


Draw or shade in the areas of your symptoms and note down any relevant history and dates below.



No Pain

0%

50%

Excruciating Pain

100%

1	Do you have any pins and needles, numbness, loss of sensation or weakness?
2	When did this problem start?
3	How did this problem start?
4	Did the pain come on immediately or did it develop over a period of hours or days?
5	Did you have any other symptoms associated with this onset? Nausea, vomiting, diarrhoea, cough, dizziness, abdominal pain, loss of bladder or bowel control, etc.
6	What did you take/do to relieve the pain? Did it help?
7	Please give a brief history of the problem. If you have already seen a doctor, specialist or therapist for this condition, give details of any examinations, tests or treatment you have had, giving dates where possible
8	What sort of movements or actions make the pain feel better or worse?
Please give any other details that you think are important or relevant.	

MEDICAL HISTORY

9	Have you had any illnesses, operations or accidents since your last treatment?				
10	Are you on any new medication since your last treatment?				
11	Have you noticed any change in your bowel or urinary habit recently? Constipation, diarrhoea, blood and/or pain on passing stool.				
12	When was the last time you had your blood pressure taken? Do you know what it was?				
13	Is your job or home life particularly stressful at the moment? If yes, do you wish to discuss this at your appointment?				
<p>Male patients answer the following questions 14,15,16 Female patients go to question 17</p>					
14	Have you noticed any change in urinary habit recently? Difficulty stopping or starting, flow rate, blood in urine, increase in frequency.				
15	Have you had a prostate examination? Give reason for the tests and the results.				
16	Have you ever noticed any pain or swelling in the testicles? If yes, did you seek medical advice? What were the results of the tests? If you have any swellings or lumps in the testicles, even if they are painless, you should consult your GP				
<h2 style="color: #e91e63;">GYNAECOLOGICAL HISTORY</h2>					
17	Have you noticed any changes in your periods since your last treatment?				
18	Do you have a coil fitted?				
19	Have you had a mammogram? If yes, how long ago? What were the results?				
20	When was your last cervical smear test?				
21	Have you had any children since your last treatment? If yes, give birth weights and any problems that there may have been with the pregnancies or births, eg: caesarean, ventouse, forceps, etc. <table style="width: 100%; border: none;"> <tr> <td style="width: 25%;">Male/Female</td> <td style="width: 25%;">Birth Weight</td> <td style="width: 25%;">Problems with labour / birth</td> <td style="width: 25%;">Age now</td> </tr> </table>	Male/Female	Birth Weight	Problems with labour / birth	Age now
Male/Female	Birth Weight	Problems with labour / birth	Age now		

The information that you provide on this form (and any other information obtained during the course of your treatment) is done so on a strictly confidential basis.

This information will be used solely for the purposes of providing osteopathic and/or any related treatment.

We will not disclose any personal information that we hold about you outside this practice without your explicit consent, except to the extent we are required or permitted by law.

The Data Protection Act ('The Act') lays down certain requirements for the protection against unauthorised disclosures of personal information.

The Act also gives you certain rights. In order that we may use your personal information for the purposes of providing osteopathic and/or any relevant treatment, we are obliged to obtain your written consent.

Accordingly, we would be obliged if you would sign the consent section below.

I consent to the use of my personal information for the purposes set out above. I also confirm that the information I have provided in this form and during the course of my treatment is, to the best of my knowledge, accurate and complete.

Signature of Patient: _____ **Date:** _____
 (If the patient is a child, then the signature of their parent or legal guardian)

If you **DO NOT** consent to your doctor being informed that, you are receiving osteopathic treatment please mark an **X** in the box.

Thank you

Before you embark upon a course of treatment with us, we need to inform you of any possible adverse side effects you might sustain from your treatment here.

Whilst such adverse effects are rare, we nevertheless need to request that you sign this document to indicate that you have been duly informed of any possible side effects of the therapies you may receive at this practice.

We have included all the therapies you can receive within this document to reduce repetition and avoid multiple consent forms.

You can give consent for just one therapy or for multiple therapies if you wish.

Parents or guardians of babies/children having Cranial Osteopathy need only read the Cranial Osteopathy section, tick the Cranial Osteopathy box and delete the others.

Whilst every attempt is made to ensure your comfort and safety, certain specific techniques may be uncomfortable and may, in some individuals, cause temporary adverse reactions.

Osteopathy

Consists of a combination of massage and manual manipulation to relieve a variety of musculo-skeletal problems.

- In patients who bruise easily (or who are currently taking medication such as steroids or Warfarin), certain massage techniques may leave some temporary bruising.
- After the first consultation, which will include a detailed physical examination, a few patients might feel an increase in symptoms due to the unaccustomed movements and actions needed to examine the spine or limb. This can last 24-48 hours and should settle with time.
- Some patients who have low blood pressure or circulation problems might experience light-headedness following a treatment; this is temporary and should settle within a few minutes/hours.
- It is not uncommon for patients to report feelings of tiredness after a treatment. We therefore advise that you avoid driving long distances or operating machinery if this occurs.
- Following some manipulations, particularly if the problem is well established or very acute, there might be a local reaction to treatment. This may last 24-48 hours, but if it does last longer, please contact the practice.

Medical Acupuncture

Acupuncture is a form of therapy in which fine needles are inserted into specific points on the body to reduce pain and muscle spasm.

Acupuncture is generally very safe. Serious side effects are very rare – less than one per 10,000 treatments.

- Drowsiness can occur after treatment in a small number of patients and, if affected, you are advised not to drive.
- Minor bleeding or bruising can occur after acupuncture (in about 3% of treatments).
- Pain during treatment can occur (in about 1% of treatments).
- Existing symptoms can get worse after treatment (in less than 3% of patients). You should tell your acupuncturist about this, but it is usually a good sign.
- Fainting can occur in certain patients, particularly at the first treatment.

Single-use, sterile, disposable needles are always used in this practice.

If you are a blood donor you will need a certificate from the practice to present at your next donor session.

Cranial Osteopathy

Cranial Osteopathy is a gentle balancing technique that is commonly used on infants and small children. Because of its gentle nature, there are very few side effects with Cranial Osteopathy

- Some babies may become either, sleepy or excited after a cranial treatment. This can last from a few hours up to a day.

Electrotherapy

Low-Level Laser Therapy (LLLT)

Low Level Laser Therapy (LLLT) reduces pain and inflammation whilst promoting repair. It can treat conditions such as osteoarthritis, sprains and strains, sports injuries, shingles, neuralgias, wounds and leg ulcers, back and neck pain.

LLLT can be safely used over metal pins, plates and in patients with pacemakers.

It cannot be used directly over a pregnant uterus, although it can safely be used over other areas of the body in a pregnant mother.

- There are normally no adverse effects from LLLT, however, occasionally, mild aching can occur after treatment. This is due to a stimulation of the local inflammation/healing process and should settle down after 24 – 48 hours.

Ultrasound

Uses ultra high frequencies of sound waves to 'micro-vibrate' soft tissues, to relax them and encourage healing. In some ways it can be thought of as a stronger version of the ultrasound machines that scan a pregnant uterus.

Ultrasound cannot be used over metal pins, plates, directly over a pacemaker or a pregnant uterus, although it can be used on other parts of the body in these cases.

- There have been some minor cases of physical pain due to "cavitation." Described as a burning feeling, cavitation is caused by the heating of the gas contained in tissue cell nuclei. This can result, temporarily, in difficulty breathing, dizziness, nausea and disorientation – although incidents of this nature are extremely rare.

Interferential

Uses electrical frequencies to stimulate peripheral nerves to either reduce pain and swelling, or stimulate blood supply and muscle contraction. It works in a similar way to a TENS machine.

It cannot be used on patients with pacemakers or over a pregnant uterus.

- There are very few known side effects with interferential, although the vacuum cups that hold the pads in place may cause superficial bruising in patients who are on blood thinning medication or steroids.
- There have been some reported cases of 'neuralgia-type' pains being temporarily worsened by interferential.

I have read the above and understand that, occasionally, there may be some unavoidable and temporary side effects resulting from my (my baby's/child's) treatment at The Osteopathic Practice. I understand that signing this form does not affect my statutory rights.

I consent to the following treatments. Please tick the therapies to which you consent.

(If you are unsure which therapies to tick, please leave the box blank and discuss this with your Osteopath.)

Osteopathy

Cranial Osteopathy

(If your baby/child is having Cranial Osteopathy, please delete the other therapies.)

Medical Acupuncture

Electrotherapy

Signed _____
(If the patient is a child, the parent or legal guardian must sign)


Date _____

If you are attending for neck or headaches, you must read and sign this before your Osteopath can treatment you.

We are obliged to inform patients requiring upper neck manipulation of any potential risks of such a manipulation before any '*High Velocity Low Amplitude Techniques*' can be used in this area. (The type of manipulation that causes a "Click" or "Popping" sound).

There have been a few occasions, following manipulation of the neck, where stroke-like symptoms and, ***in extremely rare cases***, severe injury or death has resulted.

UK Department of Health figures for 2001 indicate that this has occurred in 0.6 in a million times (six times in ten million)



A healthy vertebral artery can easily tolerate the stretching involved in an upper cervical manipulation without any ill effects.

Manipulating the lower neck does not affect the artery.

If the vertebral artery has already been damaged (Vertebral Artery Dissection or VAD) by a previous injury or, if it already has some inherited weakness, it can be at risk as it winds around the atlas (the top cervical vertebra, located just below the head).

In these cases, certain twisting motions may cause the artery to go into spasm, causing temporary dizziness and nausea, difficulty speaking or swallowing and visual disturbances.

In more severe cases, it may stretch the artery and tear the delicate or weakened lining (this causes sudden, extremely severe 'thunderclap' headaches or neck pain).

If this occurs, a thrombus (small blood clot) can then form in the injured artery wall, which can then lead to a stroke or T.I.A.

Everyday factors that have been known to injure the vertebral artery.

Ceiling painting
Nose blowing
Sneezing
Coughing

Judo
Tennis
Minor neck trauma
Yoga

Visit to the hairdressers
Reversing the car

Certain medical factors can increase your chances of damaging the vertebral artery:

High blood pressure
More common in females
Oral contraceptive use

Chronic headaches
Vascular pathology
Migraine

Smoking
Anticoagulants

Vertebral Artery Dissection tends to occur in people aged **30 – 45 years** and is rare in those over 60 years.

Before undergoing your treatment, please look through the list below and tick the appropriate box
if you have developed any of these symptoms recently or associate them with your neck pain or headache.

- Sudden onset "Thunderclap" head or neck pain. (Severe pain, unlike anything you have experienced before)
- 'Drop attacks', black outs, loss of consciousness
- Nausea, vomiting
- Dizziness, vertigo (associated with position of the head)
- Unsteady gait or feeling of weakness
- Tingling or numbness around the mouth
- Altered facial sensation
- Difficulty speaking or forming words
- Difficulty swallowing
- Tinnitus (ringing in the ears) or deafness

Have you recently been prescribed or are you currently taking any of the following?

- Anticoagulants
Eg: Warfarin
- Oral contraceptives
- Oral steroids
Eg: Prednisilone

Have you recently recovered from any of the following?

- Heart attack, stroke or TIA (transient ischaemic attack)

If you are not sure about any of these questions, please speak to your osteopath.

I have read this leaflet on the risk factors associated with cervical (neck) manipulation, and understand that there is a small (0.6 in 1,000,000) chance of vertebral artery injury following high cervical (upper neck) 'High Velocity, Low Amplitude' techniques.

Please tick the appropriate box to indicate your wishes regarding treatment:

- I agree** to have cervical (neck) 'High Velocity Low Amplitude' manipulative techniques if my course of treatment at The Osteopathic Practice requires it.
I have read the above and have had my questions regarding the potential risks answered to my satisfaction.
- I do not agree** to have cervical (neck) 'High Velocity Low Amplitude' manipulative techniques and wish to discuss possible alternatives to this treatment.
I understand that this may require more treatments or more frequent treatments if 'High Velocity Low Amplitude' techniques are not used.

Signature: _____

Date: _____

Re-Examination Questionnaire

Please complete this form in as much detail as you can. This will not only help your osteopath understand your total state of health, but can help us discover any medical conditions which may need referral to your doctor before you embark on a course of osteopathic treatment.

Your osteopath retains the right to refuse treatment if this form is not completed.

Full Name	Mr/Mrs/Miss/Ms/ other (please state) First names: Surname
Address	 Post Code
Telephone	Home Work Mobile
Email	Email address if you would like to receive information about the practice
Date of Birth	
Height	
Weight	
Occupation	
Doctor's Name	
Doctor's address	
Doctor's telephone.	